

# Disconnection between parietal and temporal areas without simultanagnosia: a case study of prosopagnosia

Manuel Alejandro Mejía, Mitchell Valdés-Sosa, Beatrice de Gelder & Maria Antonieta Bobes

**To cite this article:** Manuel Alejandro Mejía, Mitchell Valdés-Sosa, Beatrice de Gelder & Maria Antonieta Bobes (2025) Disconnection between parietal and temporal areas without simultanagnosia: a case study of prosopagnosia, *Neurocase*, 31:3, 124-132, DOI: [10.1080/13554794.2025.2489929](https://doi.org/10.1080/13554794.2025.2489929)

**To link to this article:** <https://doi.org/10.1080/13554794.2025.2489929>



Published online: 12 Apr 2025.



Submit your article to this journal [↗](#)



Article views: 157



View related articles [↗](#)



View Crossmark data [↗](#)



Citing articles: 1 View citing articles [↗](#)

RESEARCH ARTICLE



## Disconnection between parietal and temporal areas without simultanagnosia: a case study of prosopagnosia

Manuel Alejandro Mejía <sup>a,b</sup>, Mitchell Valdés-Sosa <sup>b</sup>, Beatrice de Gelder <sup>c</sup> and Maria Antonieta Bobes <sup>b</sup>

<sup>a</sup>CETYS University, Tijuana, México; <sup>b</sup>Cuban Center for Neuroscience, La Habana, Cuba; <sup>c</sup>Maastricht University, Maastricht, Netherlands

### ABSTRACT

This study presents a neuropsychological evaluation of a unique case of prosopagnosia (patient EP) with atypical lesion patterns, characterized by intact face-selective nodes but significant damage to the Vertical Occipital Fasciculus (VOF). Given the presumed interruption of ventral-parietal connectivity, we focused on assessing the potential presence of simultanagnosia and its potential relationship to his face recognition deficits. Our neuropsychological battery included tests of global and local processing, scene perception, and face recognition. Results revealed intact global processing abilities and no evidence of simultanagnosia, despite the patient's prosopagnosia. These findings suggest that EP's face recognition impairment is likely attributable to disrupted connectivity within the face processing network rather than a general deficit in global/holistic processing. This case highlights the importance of comprehensive neuropsychological assessments in atypical presentations of prosopagnosia and contributes to our understanding of the complex relationship between white matter integrity and face recognition abilities.

### ARTICLE HISTORY

Received 12 July 2024  
Accepted 1 April 2025

### KEYWORDS

Prosopagnosia;  
neuropsychological  
assessment; face recognition;  
global processing;  
simultanagnosia

### Introduction

Face recognition is a complex cognitive process that relies on the intricate interplay of various brain regions and neural networks. Acquired prosopagnosia, a selective impairment in face recognition, has traditionally been associated with lesions in face-selective areas. Two variants of prosopagnosia have been described: apperceptive, where the structural encoding of faces is impaired, and associative, where structural encoding is intact, but it is disconnected from face memory processes. Here we focus on apperceptive prosopagnosia, which has been traditionally associated with lesions in more posterior face-selective areas, such as the fusiform face area (FFA) and occipital face area (OFA) (Corrow et al., 2016), or to damage of white matter tracts, particularly the inferior longitudinal fasciculus (ILF) and inferior fronto-occipital fasciculus (IFOF), which connect nodes of the face processing circuitry (Valdés-Sosa et al., 2011). However, recent research has highlighted the importance of considering broader visual processing deficits that may contribute to face recognition impairments, particularly in cases with atypical lesion patterns (Cohen et al., 2019; Fox et al., 2008; Grill-Spector et al., 2017).

Simultanagnosia, a neurological disorder characterized by the inability to perceive multiple elements of a visual scene simultaneously, is one such deficit that can potentially impact face recognition (Meek et al., 2013; Sakurai et al., 2016). Typically, simultanagnosia is assessed by asking patients to report what they see in a complex scene where multiple objects and persons interact (e.g., the Cookie Theft Picture from the Boston Diagnostic Aphasia Examination; de Vries et al., 2022). The responses in simultanagnosia are characterized by a piece-meal approach, describing only one element at a time (e.g., the kitchen sink). Patients' responses mostly select and describe one complete object and not its parts, regardless of the size

of stimuli (with some exceptions when a specific part of an object attracts their attention initially). Patients are unaware of the rest of the scene; thus, they cannot integrate what is happening overall. This classical definition of simultanagnosia corresponds to what is also called "dorsal simultanagnosia," to distinguish it from "ventral simultanagnosia" where patients can guess the overall scene by applying the piece-meal approach effortfully (Farah, 2004). The dot-counting task has been used to distinguish between simultanagnosia types, where patients with the ventral type can count a set of black dots dispersed on a white background, while patients with the dorsal type cannot (Sakurai et al., 2016). While typically associated with bilateral parietal lobe damage, simultanagnosia can also result from lesions in other brain regions, leading to complex presentations that challenge our understanding of visual processing disorders (Cui et al., 2022).

While prosopagnosia and simultanagnosia can occur independently, their co-occurrence has been reported in various neurological conditions, particularly in some atypical lesions, and most notably, in posterior cortical atrophy (PCA) (Cui et al., 2022). PCA is a neurodegenerative syndrome characterized by progressive visual dysfunction, often presenting a combination of visual agnosias, including prosopagnosia and simultanagnosia (Cui et al., 2022). Furthermore, on the rare occasions where unilateral lesions have produced a co-occurrence of simultanagnosia and prosopagnosia, this lesion has presented on the right hemisphere (Naccache et al., 2000; Sakurai et al., 2016), leading Sakurai et al. (2016) to suggest that ventral simultanagnosia following right hemisphere damage would co-occur with prosopagnosia. Another study suggested that the face processing difficulties of their five PCA patients were produced by the reduced attentional window (i.e., simultanagnosia) after analyzing the patient's

scanning patterns in a face-matching task (Meek et al., 2013). These studies raise important questions about the potential interactions between simultanagnosia and prosopagnosia and their combined impact on visual processing.

In this study, we present a unique case of acquired prosopagnosia, patient EP, with lesions outside the typical face-selective areas, specifically the Fusiform Face Area (FFA), Occipital Face Area (OFA), Superior Temporal Sulcus (STS) and anterior Temporal Lobe (aTL) as previously reported elsewhere (Bobes et al., 2021; De Gelder et al., 2022). Additionally, the patient has damage to the white matter tracts, including the Inferior Longitudinal Fasciculus (ILF) and the Vertical Occipital Fasciculus (VOF). As a result of the VOF interruption, the patient exhibits diminished connectivity between the ventral-temporal cortex and dorsal visual areas. This disruption raises the possibility that other visual processing deficits may be contributing to the observed face recognition deficits.

The primary objective of this research is to conduct a neuropsychological evaluation to assess the potential presence of simultanagnosia in the patient EP, that could potentially underlie his face processing deficits. This assessment is crucial as it allows for a thorough examination of visual processing in a case with atypical lesion patterns and connectivity issues, helps determine whether other visual processing deficits beyond damage to face-selective areas contribute to the observed face recognition deficits, and provides an opportunity to investigate how disrupted connectivity between ventral-temporal and dorsal visual areas might impact face recognition processes.


Understanding the potential contributions of various visual processing deficits to face recognition impairments is particularly relevant in cases where prosopagnosia presents with atypical lesion patterns. By examining the visual processing abilities in a case of prosopagnosia with intact face-selective areas but disrupted white matter connectivity, we may enhance our understanding of the complex neural networks underlying visual processing and face recognition. This case study highlights the importance of comprehensive assessments in atypical presentations of prosopagnosia and underscores the need for a nuanced approach in understanding the various factors that can contribute to face recognition deficits.

## Materials and methods

### *Patient description and previous neuropsychological assessment*

Subject EP, currently 57 years old, initially presented for evaluation at age 46 following two consecutive strokes within a 1-day interval. The primary lesion was identified in the right hemisphere, affecting parts of the lingual gyrus, medial fusiform gyrus, and a small portion of the inferior temporal gyrus (see Figure 1). This lesion extended dorsally into white matter, slightly impacting the ventral part of the optic radiation. A secondary, smaller lesion was identified in the left hemisphere, confined to the middle and posterior fusiform gyrus. Further neuroimaging studies 4 and 8 years later and voxel-based morphometry (VBM) analysis consistently confirmed the location of the lesions. The structural image of EP's brain was overlaid with a region of interest mask (Bobes et al., 2021), and the functional MRI of EP in response to blocks of faces was compared to other objects (body parts, and houses) with methods described elsewhere (Van Den Stock et al., 2012). Importantly, these analyses revealed that the lesions did not directly intersect the core face areas, including the Fusiform Face Area (FFA), the Occipital Face Area (OFA), or the Superior Temporal Sulcus (STS) but did affect more medial portions of the ventral-temporal cortex within the broader face processing network (Bobes et al., 2021).

During his initial evaluation, reported previously (De Gelder et al., 2022), EP exhibited significant impairments in facial structure processing. He scored 46 out of 64 (72%) on the Facial Identity Matching Test of the FEAST Battery (De Gelder et al., 2015), where three faces are presented simultaneously, a face above in frontal view and two faces in  $\frac{3}{4}$  view below, and the patient must select which face matches the picture above. EP scored 10 out of 27 (37%; which is equivalent to a score of <25 for the Long form of the BFRT) on the short form of the original Benton Face Recognition Test (BFRT; Benton et al., 1994) where face matches have to be selected among a pool of six faces while the target face is present. As expected, EP also exhibited significant impairments in tasks that addressed face memory, scoring <50% in the Cambridge Face Memory Test (CFMT; Duchaine & Nakayama, 2006) where participants learn



**Figure 1.** T1-weighted structural image of patient EP's lesion. Arrows indicated the lesion on the right hemisphere described in the main text.

six new face identities and have to identify them in successive trials among two other distractors, and was severely impaired in recognizing famous faces, with only 5 out of 14 celebrity faces being visually recognized. EP reported relying on non-facial features like hairstyle for recognition.

EP showed intact low-level visual perception, as revealed by average scores on orientation, length, and size matching tasks from the Birmingham Object Recognition Battery. He struggled with intermediate-level tasks that are interpreted as dependent on configural information. First, he scored 10/32 on the object decision task of the BORB (De Gelder et al., 2022), where a mix of line drawings sketching real animals and non-real animals is shown and the patient had to point which were real and which were not. Second, EP struggled with the face and house part-to-whole matching tasks of the FEAST battery (De Gelder et al., 2022), where he asked to stop the task. In the part-to-whole task, a trial consists of a face (or house) displayed at the top, while two parts (eyes or nose, or window or door) are shown below, and the patient must select which is the part that corresponds to the upper image. This struggle with the part-to-whole task could be interpreted as a difficulty in configural processing for both faces and houses, but it was unclear if the reason to abort the task was a marked difficulty only for faces, for both faces and houses, or an expected difficulty and stress from the participant (this task has an average accuracy around 61–73% in middle aged control participants, Huis in 't Veld, 2015). These previous results are summarized in Table 1, including Crawford's *t* values for single case comparisons.

In further evaluations of face recognition conducted 4 and 8 years after the initial assessment, EP's performance remained impaired but showed some variability. Four years after the initial evaluation, EP scored 35 out of 54 (65%) on the BFRT and 56% on the CFMT. In the assessment conducted 8 years post-initial evaluation, EP's scored 46% on the CFMT. These results indicate a persistent deficit in face recognition abilities, consistent with the diagnosis of acquired prosopagnosia, of the apperceptive type, while also revealing subtle fluctuations in performance over time.

## Procedure

A complementary neuropsychological assessment was conducted to evaluate the potential presence of simultanagnosia

and its contribution to the patient's face recognition deficits. The assessment was performed in a quiet, well-lit room. The assessment was conducted remotely, and considered guidelines for remote neuropsychological assessment (Bilder et al., 2020). Test instructions were presented verbally and written by screen sharing. Stimuli was presented via screen sharing, except for the Navon classical task that was executed directly on the patient's computer. The patient was accompanied by a family member who was proficient in using audiovisual technology for the assessment. All researchers who participated in the remote assessment had also been part of previous in-person assessments of the patient. The patient does not wear corrective lenses for reading, he only has a prescription for larger distances.

## Neuropsychological tests

The following tests were administered to assess various aspects of visual processing, with a focus on evaluating simultanagnosia and its potential impact on face recognition. The assessment aimed to test different levels of visual integration (Piepers & Robbins, 2012). A first-order configuration was assessed with tasks of object-closure (e.g., Mooney faces and incomplete letters), then a second-order configuration was assessed with tasks of higher visual processing (e.g., face perception/memory, facial characteristics and hierarchical object recognition), and lastly, multi-object processing was assessed with complex scenes descriptions (e.g., the Cookie-Theft Picture).

- (1) **Integration of Complex Scenes.** The patient was asked to describe as accurately and thoroughly as possible what he saw that was going on in two pictures. The first picture was the "Cookie theft" from the Boston Diagnostic Aphasia Examination (Goodglass & Kaplan, 1983), and the second one was the scene from The Birthday Party test (BPT; de Vries et al., 2022). The patient was encouraged to give more information when he was not giving further responses for some time. Scoring was done following Croisile et al. (1996) guide for the first picture, and the scoring aid of the BPT. Scores were converted to Crawford's *t* values based on the norms published by de Vries et al. (2022) Complex scenes are one of the most used tests to detect simultanagnosia,

**Table 1.** Patient EP scores (points scored/maximum) in previous assessments, and conversion to T-scores based on means (SD) of controls, as reported by De Gelder et al. (2022).

Test	EP Raw score	Mean (SD) in controls	T	Interpretation
<b>Identity matching task</b>				
Upright faces	46/64	Not reported	2.9	Impaired
Inverted faces	51/64	Not reported	1.09	Normal range
<b>BFRT – Short form</b>	<25/54 <sup>a</sup>	45.4 (3.96)	–5.14	Severely impaired
<b>CFMT (%)</b>	<50/100	80.4 (11)	–2.74	Severely impaired
<b>Famous faces test</b>	5/14	Not available	Not available	Impaired
<b>BORB</b>				
Line length match	28/30	26.9 (1.6)	0.68	Normal range
Size match	26/30	27.3 (2.4)	–0.53	Normal range
Orientation match	25/30	24.8 (2.6)	0.08	Normal range
Minimal feature view	24/25	23.3 (2)	0.34	Normal range
Object decision	10/32	27 (2.2)	–7.45	Severely impaired

BFRT = Benton Face Recognition Test. BORB=Benton Object Recognition Battery. CFMT = Cambridge Face Memory Test. <sup>a</sup>Converted score from the BFRT-Short form to the Long form. T = Crawford's *t*-values to compare single cases against a control group.

characterized by an impairment in integrating the whole scene and a piece-meal approach to detect elements in the scene (de Vries et al., 2022).

- (2) **Global/Local Processing.** The patient was asked to describe as thoroughly as possible a set of 11 figures with different elements forming local and global features. The first two figures were composed of geometric figures (squares and triangles), the next two were schematic figures (cup, umbrellas, and fish, from Poirel et al., 2008), and the fifth was a classic Navon figure of a global H composed of smaller F's (Navon, 1977). The remaining figures in the set were paintings by Giuseppe Arcimboldo and Octavio Ocampo that had different figures in local and global hierarchies. This task was not time limited. Hierarchical figures, particularly those similar to Navon's, have been sensitive to simultanagnosia, where patients attention is captured by the local elements and are unable to switch to the other level (Dalrymple et al., 2007, 2013; Shenker & Roberts, 2016).

Furthermore, the patient was administered a full classical Navon task designed in PsychoPy (Peirce et al., 2019). The task consisted of four blocks (half local letters, and half attending global letters) of hierarchical letters. Each trial consisted of a hierarchical letter (e.g., a big letter "H" composed of small letters, either "H" or "S") that could be congruent (same letter on both levels) or incongruent. The patient was asked to respond which letter was represented in the local level in half of blocks, and in the global level in the other half, as fast as possible without sacrificing accuracy. There was a practice block for each level at the beginning of the task that ended when the patient had at least 80% of accuracy. At a distance of 60 cm, the size of large letters was  $3.31^\circ \times 5.23^\circ$  of visual angle, and the small letters were  $0.47^\circ \times 0.66^\circ$ . The patient remained at a constant distance from the screen throughout the task. For each trial, there was a blank screen for 500 ms, then 500 ms of fixation cross, followed by 180 ms of the hierarchical letter, that was substituted by a mask made from dots. This mask remained on-screen until the participant gave a response. The classical Navon effect consists of slower reaction times for incongruent letters only when attending the local level. The task was administered via Pavlov, and the size of stimuli was standardized by adjusting it according to the size of a credit card compared at the beginning of the task. Scores were compared to an age-matched control sample ( $n = 7$ , mean age = 52, SD age = 3.45) with a Bayesian single-case test based on Crawford's methodology (Scandola & Romano, 2021).

- (3) **Dot counting task.** The patient was asked to count the number of dots in a square, without pointing to the screen. This task was adapted from the Addenbrooke's Cognitive Examination-III (ACE-III, Hsieh et al., 2013). The dot counting task is one of the most sensitive task for ventral simultanagnosia, where patients show intact dot

counting despite the piece-meal approach used to perceive complex scenes (Sakurai et al., 2016).

- (4) **Subjective Exploration.** A semi-structured interview about day-to-day difficulties with objects or spatial abilities. Patient was asked: "Have you or anyone close to you noticed difficulties recognizing or telling apart some things in your day-to-day activities? Can be at home, work, or other activities," "How would you qualify your vision in the last 3 years? Has it improved, worsened, or stayed the same?," and "When you need to find something, either at your house or at your work or else, for example, missing keys, how do you usually manage? Do you find it easy or difficult? Have you needed help in situations like these?."
- (5) **Visual Imagery.** We adapted the Vividness of Visual Imagery Questionnaire (Blomkvist & Marks, 2023) by administering the first four items ("Think of some relative or friend whom you frequently see") and a new set of four items that more directly asked the patient to imagine the face of that same relative or friend ("Think of the face of some relative or friend whom you frequently see") asking about the clarity of the imagined face features ("overall image of the face, eyes, mouth, nose," "expressions of the face," "face smiling," and "different colors and textures of the skin"). Responses were given in a 5-point scale (5="Perfectly clear and vivid as real seeing," 1="No image at all, you only 'know' that you are thinking of the object"), in two conditions: eyes open and eyes closed. Scores were converted to Crawford's  $t$  values based on the norms published by McKelvie (1995) for individual items.
- (6) **Object Closure.** The first task was the incomplete letters from the ACE-III (Hsieh et al., 2013), where the patient is asked to identify letters that are missing some parts. The second task was an adaptation of the Mooney Face Test (Mooney, 1957). The patient was asked to select which of three images shown in a slide was a face. Stimuli remained on the screen for a maximum of 3 s. There were 20 items and a practice trial. Stimuli was one Mooney face per trial and two distractors (scrambled Mooney stimuli) selected from the set published by Schwiedrzik et al. (2018).
- (7) **Facial Characteristics Task.** The patient was asked to classify a set of 20 neutral faces by their age (young vs. old) and gender (male vs. female) (Tranel et al., 1988). There were 20 photos of males and 20 photos of females. Half of each set was young (19–31 years old) and half older (69–80 years old). Stimuli remained on the screen for a maximum of 3 s. Images were selected from the FACES database (Ebner et al., 2010).

### Data analysis

Performance on each task was evaluated against data from controls described for each task in Methods, by estimating Crawford's  $T$  values for single-case comparisons, and impairments were based on one-tail tests at 0.05 alpha level. Qualitative analysis of response patterns was conducted to



identify signs of simultanagnosia, such as piecemeal processing, difficulty with global integration, and impaired perception of multiple elements simultaneously. In the classical Navon task, we estimated accuracy for all trials (excluding practice trials), response times averages were estimated only on correct trials and response times above 200 ms, and a Bayesian multilevel single-case analysis (Scandola & Romano, 2021) was used to estimate the significance of congruency (congruent vs incongruent trials), condition (global vs local blocks) and their interaction. This multilevel analysis allows the test of those effects in the patient and the difference of each effect to the control group. Bayes factors ( $BF_{10}$ ) above 3 indicate evidence in favor of the alternative hypotheses, and  $BF_{10} < 1/3$ , in favor of the null hypotheses.

### Ethical considerations

The study was conducted in accordance with the Declaration of Helsinki and approved by the institutional ethics committee. Informed consent was obtained from the patient prior to the assessment.

## Results

The patient underwent a comprehensive neuropsychological assessment to evaluate potential simultanagnosia and its contribution to face recognition deficits. The results are described below and summarized in Table 2.

### Integration of complex scenes and hierarchical figures

For the Cookie Theft picture, the patient's performance was within the normal range compared to healthy subjects (de Vries et al., 2022), scoring 18 out of 22 overall (Mean healthy subjects = 18.61, SD = 2.82). In the Birthday Party Test (BPT), the patient performed well, scoring 28 out of 40 total (Mean healthy subjects = 26.49, SD = 5.71) (de Vries et al., 2022). Notably, the patient identified all persons/animals and performed above average in actions/relations (see Table 2).

In the Hierarchical Figures test, the patient successfully reported both global and local levels for all 11 figures, although they took longer to recognize the global level in the 11th figure.

In the classical Navon task, the patient had good accuracy in general (86.9%), with lower accuracy for local letters (see Table 2) which indicates a global advantage as found in

**Table 2.** Patient EP scores in the battery (points scored/maximum), and conversion to T-scores based on means (SD) of controls.

Test	EP Raw score	Mean (SD) in controls	T	Interpretation
<b>BFRT – Long form</b>	35/54	45.4 (3.96)	−2.63	Severely impaired
<b>Famous faces test</b>	36/73	NA	NA	Impaired
<b>CFMT (%)</b>				
4 years after lesion	56/100	80.4 (11)	−2.22	Severely impaired
8 years after lesion	46/100	80.4 (11)	−3.13	Severely impaired
<b>Hierarchical figures</b>	11/11	At ceiling	NA	Normal range
<b>Cookie theft</b>				
Subjects	3/4	3.48 (0.56)	−0.86	Normal range
Objects	8/11	8.38 (2.02)	−0.19	Normal range
Places	2/2	1.69 (0.57)	0.54	Normal range
Actions/facts	5/7	5.06 (0.97)	−0.06	Normal range
Total	18/23	18.61 (2.82)	−0.22	Normal range
<b>The Birthday Party (TBP)</b>				
Persons/animal	9/9	7.78 (1.52)	0.80	Normal range
Objects	11/18	11.42 (3.99)	−0.11	Normal range
Actions/relations	8/13	7.29 (1.94)	0.37	Normal range
Total	28/40	26.49 (5.71)	0.26	Normal range
<b>ACE-III</b>				
Dot counting	4/4	At ceiling	NA	Normal range
Incomplete letters	4/4	At ceiling	NA	Normal range
<b>Mooney faces</b>	19/20	Near ceiling	NA	Normal range
<b>Facial characteristics</b>				
Categorize age	20/20	At ceiling	NA	Normal range
Categorize sex	19/20	Near ceiling	NA	Normal range
<b>Navon, Accuracy (% correct)</b>				
Global, consistent	98	98 (2.1)	−0.09	Normal range
Global, inconsistent	94	97 (3.3)	−0.85	Normal range
Local, consistent	89	97 (3.2)	−2.32	Impaired
Local, inconsistent	68	87 (18.1)	−0.99	Normal range
<b>Navon, RT (milliseconds)</b>				
Global, consistent	582	569 (136)	0.09	Normal range
Global, inconsistent	585	559 (132)	0.18	Normal range
Local, consistent	812	662 (134)	1.05	Normal range
Local, inconsistent	845	784 (194)	0.29	Normal range
<b>Inverse Efficiency Scores (IES)</b>				
Global, consistent	597	582 (139)	0.10	Normal range
Global, inconsistent	624	577 (141)	0.31	Normal range
Local, consistent	914	684 (151)	1.43	Normal range
Local, inconsistent	1252	905 (287)	1.13	Normal range
<b>Global-Local bias</b>	0.704	0.781 (.076)	−0.95	Normal range

NA = Not available. BFRT = Benton Face Recognition Test. CFMT = Cambridge Face Memory Test. ACE-III=Addenbrooke's Cognitive Examination-III. T=Crawford's t-values to compare single cases against a control group. RT=Response time.

controls. The reaction times (RT) on correct trials showed the classical congruency effect during the local attention condition confirmed with a significant interaction between congruency and condition in a Bayesian multilevel single-case analysis ( $BF_{10} = 5.7 \times 10^6$ ), and no significant difference between patient and controls ( $BF_{10} = 0.95$ ). The congruency effect during local attention was small (incongruent trials were 33 ms slower on average), but EP showed a speed-accuracy trade-off by sacrificing accuracy for incongruent trials. To control for this trade-off, we estimated Inverse Efficiency Scores ( $IES = RT/accuracy$ ) for each condition which further confirmed a global advantage in EP with lower efficiency with local letters, and particularly with incongruent local letters (see Table 2). Lastly, EP showed a global bias similar to controls in a Global-Local bias index used by Duchaine et al. (2007) that estimates the ratio of RT for global/local conditions overall.

### Detection of multiple objects

The patient achieved a perfect score (4/4) on the dot counting task from the ACE-III.

### Subjective exploration

The patient reported that neither him nor other people have noticed other difficulties of visual recognition apart from their face recognition difficulties and the upper-left scotoma. Patient reports not having any change in their vision in the last 3 years, and does not report any further difficulties with vision, visual search (e.g., finding misplaced keys) or spatial navigation in their work, which involves high visuospatial skills.

### Visual imagery

On the Vividness of Visual Imagery Questionnaire (VVIQ), focusing on facial imagery, the patient's scores ranged from 3 to 5 indicating moderate to high vividness of visual imagery for facial features and associated details. There was no difference between conditions with eyes closed (mean = 4.1) and eyes open (mean = 4.1), and no difference in any item in comparison to controls (see Table 3). The lowest rated item was when the patient was asked to imagine the "overall face, eyes, mouth and nose" of the person, which the patient rated as 3 ("Moderately clear and lively").

### Object closure

The patient scored perfectly (4/4) on the Incomplete Letters task from the ACE-III. The patient demonstrated excellent performance on the Mooney Faces test, correctly identifying 19 out of 20 items. The error was made on item 17, but the patient self-corrected. This item had a difficulty higher than the average published stimuli, with a 72% of correct responses, compared to the average of 88.5% obtained by non-neurological participants (Schwiedrzik et al., 2018).

### Facial characteristics task

In the facial characteristics classification task, the patient scored perfectly (20/20) for age discrimination and nearly perfectly (19/20) for gender discrimination, with one initial error that was self-corrected.

### Clinical judgment

Based on the Birthday Party Test performance, our qualitative analyses concluded that: 1) the participant described the picture accurately based on both details and the whole, 2) some encouragement was needed regarding the whole picture description, in particular about the type of scene that it was (i.e., a birthday party), 3) the encouragement was very helpful in improving the description, and 4) the possibility of simultanagnosia was deemed "very unlikely."

Overall, the patient's performance across these tests suggests intact global processing abilities and no clear evidence of simultanagnosia. The patient demonstrated good performance in tasks requiring integration of complex scenes, object closure and recognition, and assessment of facial characteristics. The patient also showed the classical congruency effect while attending the local level of hierarchical letters, but not during attention to the global level. While some initial difficulties were noted in describing the whole scene in the BPT, these were readily overcome with minimal encouragement.

### Discussion

This study aimed to evaluate the potential presence of simultanagnosia in a prosopagnosic patient with atypical lesion patterns and disrupted connectivity between visual and parietal areas. The complementary neuropsychological assessment revealed several key findings that shed light on the patient's

**Table 3.** Patient EP scores in the vividness of visual imagery questionnaire (VVIQ, points scored in a scale from 1 to 5), and conversion to T-scores based on means (SD) of controls.

VVIQ		Eyes open	Eyes closed	EP's Mean	T	Interpretation
Face of relative or friend	Overall face, eyes, mouth, nose.	3	3	3	−0.99	Normal range
	Expressions of the face.	4	4	4	0.44	Normal range
	Face smiling.	4	5	4.5	1.15	Normal range
	Colors and textures of the skin.	4	3	3.5	−0.28	Normal range
Relative or friend	Contour of face, head, shoulders, body.	4	4	4	0.44	Normal range
	Poses of head, attitudes of the body.	4	4	4	0.44	Normal range
	Carriage, length of step, in walking.	5	5	5	1.87	Normal range
	Colors worn in familiar clothes.	5	5	5	1.87	Normal range

T = Crawford's t-values to compare single cases against a control group, obtained by averaging the scores for eyes open and closed and compared to a mean of 3.693 and SD = 0.692 based on a meta-analysis of normative scores (McKelvie, 1995).

visual processing abilities and the nature of their face recognition deficits.

Contrary to our initial hypothesis, the results strongly suggest that simultanagnosia is not a significant contributing factor to this patient's prosopagnosia. The patient demonstrated intact configural processing abilities across various tasks with varying levels of integration. EP showed intact first-order configuration as shown in the Mooney task, despite his deficit in the second-order configuration for faces as suggested by the tasks of face perception and memory. Also, EP showed intact integration of complex scenes and hierarchical figure recognition. This finding is crucial, as it indicates that the face recognition deficits observed in this case are likely not due to a general impairment in processing multiple elements of a visual scene simultaneously, which is characteristic of simultanagnosia (Dalrymple et al., 2013).

The patient's performance on the Birthday Party Test (BPT) and the Cookie Theft picture was particularly informative. Despite initial hesitation in describing the whole scene in the BPT, the patient was able to overcome this with minimal encouragement, ultimately performing at or above the level of healthy controls. This suggests that while there might be a slight tendency toward local processing, the patient retains the ability to integrate information globally when prompted. This pattern is inconsistent with the severe global processing deficits typically seen in simultanagnosia (Dalrymple et al., 2013).

The excellent performance on tasks requiring assessment of facial characteristics, such as age and gender discrimination, is intriguing given the patient's prosopagnosia. This dissociation between intact facial characteristics assessment and impaired face recognition supports the idea that prosopagnosia in this case may be more related to higher-level integration of facial features or accessing facial memories, rather than a fundamental deficit in perceiving facial elements (Barton & Corrow, 2016).

The patient's ability to recognize Mooney faces, which require holistic processing in a first-order configuration (Piepers & Robbins, 2012) due to their high-contrast, two-tone nature, further argues against a global processing deficit. This finding aligns with research suggesting that some prosopagnosic patients retain the ability to perceive the global form of faces while still struggling with individual face recognition (Rossion, 2008).

The intact performance on visual imagery tasks, as assessed by the VVIQ, is noteworthy. Patients with posterior cortical atrophy (PCA) commonly show both deficits in visual imagery (Dietz et al., 2023) and simultanagnosia (Maia Da Silva et al., 2017). Nevertheless, lesions that cause simultanagnosia may co-occur with certain types of visual imagery deficits but not others, e.g., deficits in imagery for mental rotation or spatial processing, but not in imagery for color and shape (Foley et al., 2020). Our patient showed neither type of impairment, narrowing their deficit to a highly specific impairment for face perception, with possibly spared visual imagery for faces. This dissociation between face perception and face imagery further supports the complexity of face processing networks and the potential for selective impairments within this system.

Given these findings, we must consider the role of white matter tract damage in our patient's prosopagnosia. The lesions to both the ILF (23% of volume) and VOF (60% of volume) likely disrupt the connectivity within the face processing network (Bobes et al., 2021). While the ILF connects occipital visual areas with temporal regions and high-level visual areas within the occipito-temporal region, the VOF damage may impair communication between ventral and dorsal visual streams (Yeatman et al., 2014). This pattern of disconnection could explain the apperceptive nature of our patient's prosopagnosia, disrupting the integration of facial features and configural face processing.

The disrupted connectivity between visual and parietal areas observed in this patient might lead to face recognition deficits even in the absence of damage to core face-selective regions like the FFA and OFA (Alegret et al., 2009). Kay & Yeatman (2017) suggested that parietal areas may modulate the size of perceptual receptive fields when attending to faces in healthy subjects via VOF. We put forward the hypothesis that the patient's deficit with faces may stem from impaired modulation of spatial processing specific to faces, rather than a general deficit in global visual processing.

The absence of simultanagnosia symptoms in our patient, despite the ventral-parietal disconnection caused by damage to the vertical occipital fasciculus (VOF), can be explained by several factors. One possibility is that simultanagnosia typically requires more extensive or bilateral damage to ventral-parietal connections, and the unilateral lesion in our patient may not be sufficient to produce detectable symptoms. This is supported by research showing that simultanagnosia is often associated with bilateral parietal lesions or diffuse cortical atrophy (Chechlacz et al., 2012). Furthermore, the preservation of other white matter pathways or cortical areas involved in visual attention and spatial integration could be mitigating the effects of the VOF disconnection. For instance, the intact superior longitudinal fasciculus (SLF) might be compensating for the VOF damage, as the SLF is known to play a crucial role in spatial attention and visual processing (de Schotten et al., 2011).

This dissociation between prosopagnosia and the absence of simultanagnosia underscores the complexity of neural networks involved in visual processing and highlights the need for further research on the relationship between brain connectivity and specific visual deficits. It suggests that the brain's visual processing system may have redundant pathways or compensatory mechanisms that can maintain global visual integration even in the face of specific white matter tract damage.

This case highlights the complexity of face recognition processes and the importance of considering network-level disruptions in addition to focal lesions when studying prosopagnosia. It also underscores the value of comprehensive neuropsychological assessments in differentiating between various visual processing deficits and understanding the specific nature of face recognition impairments in atypical cases.

Future research could benefit from combining detailed behavioral assessments with advanced neuroimaging techniques to further elucidate the relationship between structural connectivity, functional activation patterns, and face recognition abilities in prosopagnosic patients with atypical lesion patterns.



In conclusion, while our patient presents with apperceptive prosopagnosia, the neuropsychological profile does not support the presence of simultanagnosia. The face recognition deficits in this case are more likely attributable to disrupted connectivity within the face processing network rather than a general impairment in global visual processing. This study contributes to our understanding of the diverse presentations of prosopagnosia and emphasizes the need for nuanced approaches in diagnosing and understanding face recognition disorders.

## Acknowledgements

We are thankful to EP and his wife for their generous participation in the assessments for this study. During the preparation of this work, the authors used ChatGPT-4o in order to improve readability and shorten paragraphs in the Introduction and Discussion. After using these tools, the authors reviewed and edited the content as needed and take full responsibility for the content of the publication.

## Disclosure statement

No potential conflict of interest was reported by the author(s).

## Funding

MM received funding from the Institutional Research Coordination [Project #312] of CETYS University. MVS, and MAB were supported by project PN305LH013-061 of the National Program of Neuroscience and Neurotechnology of the Ministry of Science, Technology, and Environment of Cuba. BdG was supported by the European Research Council ERC Synergy grant (Grant agreement 856495, Relevance), by the European Union's Horizon 2020 research and innovation program (Grant agreement 101017884, GuestXR), the European Union's Horizon Europe research and by the innovation program (Grant agreement 101070278, ReSilence). The funding sources had no further involvement in the design, data collection, analysis, or interpretation of the data for this study.

## ORCID

Manuel Alejandro Mejía  <http://orcid.org/0000-0002-6049-4941>

Mitchell Valdés-Sosa  <http://orcid.org/0000-0002-1580-8374>

Beatrice de Gelder  <http://orcid.org/0000-0003-1667-8986>

Maria Antonieta Bobes  <http://orcid.org/0000-0003-0024-3881>

## Data availability statement

The data that support the findings of this study are available either within the article (Tables 1–3), openly available in the OpenScienceFramework at <https://osf.io/zkt7h/>, and other restricted data are available on request from the corresponding author, MM.

## References

- Alegret, M., Boada-Rovira, M., Vinyes-Junqué, G., Valero, S., Espinosa, A., Hernández, I., Modinos, G., Rosende Roca, M., Mauleón, A., Becker, J. T., & Tárraga, L. (2009). Detection of visuo-perceptual deficits in preclinical and mild Alzheimer's disease. *Journal of Clinical and Experimental Neuropsychology*, 31(7), 860–867. <https://doi.org/10.1080/13803390802595568>
- Barton, J. J. S., & Corrow, S. L. (2016). Recognizing and identifying people: A neuropsychological review. *Cortex, A Journal Devoted to the Study of the Nervous System and Behavior*, 75, 132–150. <https://doi.org/10.1016/j.cortex.2015.11.023>
- Benton, A., Sivan, A. B., Hamsher de, K. S., Varney, N. R., & Spreen, O. (1994). *Contributions to neuropsychological assessment: A clinical manual* (2nd ed.). Oxford University Press.
- Bilder, R. M., Postal, K. S., Barisa, M., Aase, D. M., Cullum, C. M., Gillaspay, S. R., Harder, L., Kanter, G., Lanca, M., Lechuga, D. M., Morgan, J. M., Most, R., Puente, A. E., Salinas, C. M., & Woodhouse, J. (2020). Inter organizational practice committee recommendations/guidance for teleneuropsychology in response to the COVID-19 Pandemic. *Archives of Clinical Neuropsychology*, 35(6), 647–659. <https://doi.org/10.1093/arclin/acia046>
- Blomkvist, A., & Marks, D. F. (2023). Defining and 'diagnosing' aphantasia: Condition or individual difference? *Cortex, A Journal Devoted to the Study of the Nervous System and Behavior*, 169, 220–234. <https://doi.org/10.1016/j.cortex.2023.09.004>
- Bobes, M. A., Van den Stock, J., Zhan, M., Valdes-Sosa, M., & de Gelder, B. (2021). Looking beyond indirect lesion network mapping of prosopagnosia: Direct measures required. *Brain*, 144(9), e75–e75. <https://doi.org/10.1093/brain/awab276>
- Chechacz, M., Rotshtein, P., Hansen, P. C., Riddoch, J. M., Deb, S., & Humphreys, G. W. (2012). The neural underpinnings of simultanagnosia: Disconnecting the visuospatial attention network. *Journal of Cognitive Neuroscience*, 24(3), 718–735. [https://doi.org/10.1162/jocn\\_a\\_00159](https://doi.org/10.1162/jocn_a_00159)
- Cohen, A. L., Soussand, L., Corrow, S. L., Martinaud, O., Barton, J. J. S., & Fox, M. D. (2019). Looking beyond the face area: Lesion network mapping of prosopagnosia. *Brain*, 142(12), 3975–3990. <https://doi.org/10.1093/brain/awz332>
- Corrow, S. L., Dalrymple, K. A., & Barton, J. J. S. (2016). Prosopagnosia: Current perspectives. *Eye and Brain*, 8, 165–175. <https://doi.org/10.2147/EB.S92838>
- Croisile, B., Ska, B., Brabant, M.-J., Duchene, A., Lepage, Y., Aimard, G., & Trillet, M. (1996). Comparative study of oral and written picture description in patients with Alzheimer's disease. *Brain and Language*, 53(1), 1–19. <https://doi.org/10.1006/brln.1996.0033>
- Cui, Y., Liu, Y., Yang, C., Cui, C., Jing, D., Zhang, X., Chen, Y., Li, B., Liang, Z., Chen, K., Zhang, Z., & Wu, L. (2022). Brain structural and functional anomalies associated with simultanagnosia in patients with posterior cortical atrophy. *Brain Imaging and Behavior*, 16(3), 1148–1162. <https://doi.org/10.1007/s11682-021-00568-8>
- Dalrymple, K. A., Barton, J. J. S., & Kingstone, A. (2013). A world unglued: Simultanagnosia as a spatial restriction of attention. *Frontiers in Human Neuroscience*, 7. <https://doi.org/10.3389/fnhum.2013.00145>
- Dalrymple, K. A., Kingstone, A., & Barton, J. J. S. (2007). Seeing trees or seeing forests in simultanagnosia: Attentional capture can be local or global. *Neuropsychologia*, 45(4), 871–875. <https://doi.org/10.1016/j.neuropsychologia.2006.07.013>
- De Gelder, B., Huis in 't Veld, E. M. J., & Van Den Stock, J. (2015). The facial expressive action stimulus test. A test battery for the assessment of face memory, face and object perception, configuration processing, and facial expression recognition. *Frontiers in Psychology*, 6. <https://doi.org/10.3389/fpsyg.2015.01609>
- De Gelder, B., Huis in 't Veldt, E., Zhan, M., & Van Den Stock, J. (2022). Acquired prosopagnosia with structurally intact and functional fusiform face area and with face identity-specific configuration processing deficits. *Cerebral Cortex*, 32(21), 4671–4683. <https://doi.org/10.1093/cor/bhab509>
- de Schotten, M. T., Dell'acqua, F., Forkel, S. J., Simmons, A., Vergani, F., Murphy, D. G. M., & Catani, M. (2011). A lateralized brain network for visuospatial attention. *Nature Neuroscience*, 14(10), 1245–1246. <https://doi.org/10.1038/nn.2905>
- de Vries, S. M., Tucha, O., Melis-Dankers, B. J. M., Vrijling, A. C. L., Ribbers, S., Cornelissen, F. W., & Heutink, J. (2022). The birthday party test (BPT): A new picture description test to support the assessment of simultanagnosia in patients with acquired brain injury. *Applied Neuropsychology Adult*, 29(3), 383–396. <https://doi.org/10.1080/23279095.2020.1763998>
- Dietz, C. D., Albonico, A., Tree, J. J., & Barton, J. J. S. (2023). Visual imagery deficits in posterior cortical atrophy. *Cognitive Neuropsychology*, 40(7–8), 351–366. <https://doi.org/10.1080/02643294.2024.2346362>
- Duchaine, B., Germine, L., & Nakayama, K. (2007). Family resemblance: Ten family members with prosopagnosia and within-class object agnosia. *Cognitive Neuropsychology*, 24(4), 419–430. <https://doi.org/10.1080/02643290701380491>

- Duchaine, B., & Nakayama, K. (2006). The cambridge face memory test: Results for neurologically intact individuals and an investigation of its validity using inverted face stimuli and prosopagnosic participants. *Neuropsychologia*, 44(4), 576–585. <https://doi.org/10.1016/j.neuropsychologia.2005.07.001>
- Ebner, N. C., Riediger, M., & Lindenberger, U. (2010). FACES-A database of facial expressions in young, middle-aged, and older women and men: Development and validation. *Behavior Research Methods*, 42(1), 351–362. <https://doi.org/10.3758/BRM.42.1.351>
- Farah, M. J. (2004). *Visual agnosia* (2nd ed.). MIT.
- Foley, J. A., Bayraktar, I., Hyare, H., & Caine, D. (2020). Impaired spatial processing in visual perception, imagery and art-making following parieto-occipital infarcts. *Cortex, A Journal Devoted to the Study of the Nervous System and Behavior*, 126, 355–367. <https://doi.org/10.1016/j.cortex.2020.01.023>
- Fox, C. J., Iaria, G., & Barton, J. S. (2008). Disconnection in prosopagnosia and face processing. *Cortex, A Journal Devoted to the Study of the Nervous System and Behavior*, 44(8), 996–1009. <https://doi.org/10.1016/j.cortex.2008.04.003>
- Goodglass, H., & Kaplan, E. (1983). *The assessment of aphasia and related disorders* (2nd ed.). Lea & Febiger.
- Grill-Spector, K., Kay, K., & Weiner, K. S. (2017). The functional neuroanatomy of face processing: Insights from neuroimaging and implications for deep learning. In B. Bhanu & A. Kumar (Eds.), *Deep learning for biometrics* (pp. 3–31). Springer International Publishing. [https://doi.org/10.1007/978-3-319-61657-5\\_1](https://doi.org/10.1007/978-3-319-61657-5_1)
- Hsieh, S., Schubert, S., Hoon, C., Mioshi, E., & Hodges, J. R. (2013). Validation of the Addenbrooke's cognitive examination III in frontotemporal dementia and Alzheimer's disease. *Dementia and Geriatric Cognitive Disorders*, 36(3–4), 242–250. <https://doi.org/10.1159/000351671>
- Huis in 't Veld, E. (2015). *From individual to crowd perception: How motions and emotions influence the perception of identity, social interactions, and bodily muscle activations* [Doctoral Dissertation]. Tilburg University.
- Kay, K. N., & Yeatman, J. D. (2017). Bottom-up and top-down computations in word- and face-selective cortex. *eLife*, 6, e22341. <https://doi.org/10.7554/eLife.22341>
- Maia Da Silva, M. N., Millington, R. S., Bridge, H., James-Galton, M., & Plant, G. T. (2017). Visual dysfunction in posterior cortical atrophy. *Frontiers in Neurology*, 8, 389. <https://doi.org/10.3389/fneur.2017.00389>
- McKelvie, S. J. (1995). The VVIQ as a psychometric test of individual differences in visual imagery vividness: A critical quantitative review and plea for direction. *Journal of Mental Imagery*, 19(3–4), 1–106.
- Meek, B. P., Locheed, K., Lawrence-Dewar, J. M., Shelton, P., & Marotta, J. J. (2013). Posterior cortical atrophy: An investigation of scan paths generated during face matching tasks. *Frontiers in Human Neuroscience*, 7. <https://doi.org/10.3389/fnhum.2013.00309>
- Mooney, C. M. (1957). Age in the development of closure ability in children. *Canadian Journal of Psychology / Revue Canadienne de Psychologie*, 11(4), 219–226. <https://doi.org/10.1037/h0083717>
- Naccache, L., Slachevsky, A., Levy, R., & Dubois, B. (2000). Simultanagnosia in a patient with right brain lesions. *Journal of Neurology*, 247(8), 650–651. <https://doi.org/10.1007/s004150070138>
- Navon, D. (1977). Forest before trees: The precedence of global features in visual perception. *Cognitive Psychology*, 9(3), 353–383. [https://doi.org/10.1016/0010-0285\(77\)90012-3](https://doi.org/10.1016/0010-0285(77)90012-3)
- Peirce, J., Gray, J. R., Simpson, S., MacAskill, M., Höchenberger, R., Sogo, H., Kastman, E., & Lindeløv, J. K. (2019). PsychoPy2: Experiments in behavior made easy. *Behavior Research Methods*, 51(1), 195–203. <https://doi.org/10.3758/s13428-018-01193-y>
- Piepers, D. W., & Robbins, R. A. (2012). A review and clarification of the terms “holistic. Configural,” and “Relational” in the Face Perception Literature. *Frontiers in Psychology*, 3. <https://doi.org/10.3389/fpsyg.2012.00559>
- Poirel, N., Pineau, A., & Mellet, E. (2008). What does the nature of the stimuli tell us about the global precedence effect?. *Acta Psychologica*, 127(1), 1–11. <https://doi.org/10.1016/j.actpsy.2006.12.001>
- Rossion, B. (2008). Picture-plane inversion leads to qualitative changes of face perception. *Acta Psychologica*, 128(2), 274–289. <https://doi.org/10.1016/j.actpsy.2008.02.003>
- Sakurai, Y., Hamada, K., Tsugawa, N., & Sugimoto, I. (2016). Ventral simultanagnosia and prosopagnosia for unfamiliar faces due to a right posterior superior temporal sulcus and angular gyrus lesion. *Neurocase*, 22(1), 122–129. <https://doi.org/10.1080/13554794.2015.1066827>
- Scandola, M., & Romano, D. (2021). Bayesian multilevel single case models using ‘stan’. A new tool to study single cases in neuropsychology. *Neuropsychologia*, 156, 107834. <https://doi.org/10.1016/j.neuropsychologia.2021.107834>
- Schwiedrzik, C. M., Melloni, L., & Schurger, A. (2018). Mooney face stimuli for visual perception research. *PLOS ONE*, 13(7), e0200106. <https://doi.org/10.1371/journal.pone.0200106>
- Shenker, J. I., & Roberts, M. H. (2016). Simultanagnosia: When all you can see are trees, the forest still rules. *Neurocase*, 22(3), 289–293. <https://doi.org/10.1080/13554794.2015.1137949>
- Tranel, D., Damasio, A. R., & Damasio, H. (1988). Intact recognition of facial expression, gender, and age in patients with impaired recognition of face identity. *Neurology*, 38(5), 690–690. <https://doi.org/10.1212/WNL.38.5.690>
- Valdés-Sosa, M., Bobes, M. A., Quiñones, I., García, L., Valdes-Hernandez, P. A., Iturria, Y., Melie-Garcia, L., Lopera, F., & Asencio, J. (2011). Covert face recognition without the fusiform-temporal pathways. *Neuroimage*, 57(3), 1162–1176. <https://doi.org/10.1016/j.neuroimage.2011.04.057>
- Van Den Stock, J., De Gelder, B., De Winter, F.-L., Van Laere, K., & Vandenbulcke, M. (2012). A strange face in the mirror. Face-selective self-misidentification in a patient with right lateralized occipito-temporal hypo-metabolism. *Cortex, A Journal Devoted to the Study of the Nervous System and Behavior*, 48(8), 1088–1090. <https://doi.org/10.1016/j.cortex.2012.03.003>
- Yeatman, J. D., Weiner, K. S., Pestilli, F., Rokem, A., Mezer, A., & Wandell, B. A. (2014). The vertical occipital fasciculus: A century of controversy resolved by in vivo measurements. *Proceedings of the National Academy of Sciences*, 111(48). <https://doi.org/10.1073/pnas.1418503111>